

Disordered Critical Wave functions in Random Bond Models in Two Dimensions — Random Lattice Fermions at $E = 0$ without Doubling —

Yasuhiro Hatsugai^{1*}, Xiao-Gang Wen², and Mahito Kohmoto³

¹ *Department of Applied Physics, University of Tokyo, 7-3-1 Hongo Bunkyo-ku, Tokyo 113, Japan*

² *Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139*

³ *Institute for Solid State Physics, University of Tokyo, 7-22-1 Roppongi Minato-ku Tokyo 106, Japan*
(March 14, 1996)

Random bond Hamiltonians of the π flux state on the square lattice are investigated. It has a special symmetry and all states are paired except the ones with zero energy. Because of this, there are always zero-energy states (or zero-modes). The states near the zero-modes are described by massless Dirac fermions. For the zero-mode in a system with odd numbers of sites, we can construct a random lattice fermion without a doubling and quite large systems (up to 801×801) are treated numerically. We clearly demonstrate that the zero-mode is given by a critical wave function. For the multifractal behavior of the wave function, the agreement with the effective field theory is demonstrated.

PACS numbers: 73.40H, 61-43-j, 72-15R

Random critical points in *two dimensions* appear in quantum Hall systems and in systems with spin-orbit coupling. Both critical points have important experimental implications, and both are in strong coupling limit. Because of that we are still unable to calculate their critical exponents analytically, despite the efforts of more than a decade. Recently, a class of exactly solvable two-dimensional random critical points was found. [1,2] Those critical points appear in models of Dirac fermions coupled to random gauge fields. However, it was pointed out that those critical points contain infinite number of relevant directions. Thus it seems that it is almost impossible to realize those random critical points in a lattice model.

In this paper, we study random bond models on the square lattice with only nearest neighbor hopping. The models have π flux per plaquette on the average. When all hopping matrix elements are real (in this case the model will be called random hopping (RH) model), we find that with weak randomness and in a continuum approximation, the states near zero energy are described by two copies of Dirac fermions coupled to *imaginary* random chemical potential, real random mass, and *imaginary* gauge fields. The results in Ref. [1,3], suggest that the continuum model has a line of random critical points, and due to the *imaginary* gauge fields and the *imaginary* chemical potential, the critical points has only one marginally relevant direction if we fix the energy to be zero. Thus it could be possible to fine tune parameters to reach the critical line. In fact, the zero-energy states of the fine tuned model are found to have a multifractal behavior and are critical. [4] We also present direct numerical results indicating that the zero-energy states in the generic random bond model also appear to be critical with multifractal behaviors at least for the systems of sizes up to 801×801 . The multifractal scaling function

$f(\alpha)$ for the fine-tuned RH model is calculated numerically which agrees with the exact result of the continuum theory. [1,2]. Another interesting point in our treatment is that we can avoid the fermion doubling as far as the zero-energy states are concerned.

In one dimension, some of the wave functions of the quasiperiodic systems show the critical behavior clearly. [5] In two dimensions, it was difficult to show the criticality clearly due to the limitation of available system sizes. However, since our models have several specialties as described below, we can treat quite large two dimensional systems up to 801×801 which enables a clear demonstration.

The wave functions in a two-dimensional random system are believed to be always localized if the system has the time reversal symmetry. Thus it is quite interesting that our models have random critical points. There are several reasons why the critical states are allowed in our models in two dimensions. Our Random bond model has a very special property that the random Hamiltonian anti-commute with the operator γ defined below. This means that the eigenstates always appear in pairs with energies E and $-E$. Therefore the states near zero energy are quite special, and one may wonder if they are critical states or not. We may also consider a pure system with a small diagonal hopping m (which break the symmetry mentioned above). [6] The symmetry is recovered by taking a limit $m \rightarrow 0$. When $m \neq 0$, there is an energy gap near $E = 0$ and there are two energy bands. The Hall conductance of the lower band is $+\text{sign}(m)$ and $-\text{sign}(m)$ for the higher one. Since the nonzero Hall conductance is carried by extended states, there are paired extended states in the lower and the higher bands. By taking $m \rightarrow 0$, the paired extended states merges and disappear at $E = 0$. It suggests the criticality of the

$E = 0$ states. Also from a point of the perturbation theory, the density of states of the pure state vanishes at zero energy. Therefore a usual treatment of the perturbative consideration has to be modified. This property, we believe, is also responsible for the appearance of the critical states at zero energy.

Our Hamiltonian is

$$H = \sum_{\langle i,j \rangle} c_i^\dagger t_{ij} c_j + h.c. \quad (1)$$

where the summation is over the nearest-neighbor bonds. We shall analyze the following two types of random bond models near the π flux: (i) *random gauge model* $t_{j+\hat{x},j} = (-)^{j_y} e^{i\delta\theta_x(j)}$, $t_{j+\hat{y},j} = e^{i\delta\theta_y(j)}$, (ii) *random hopping (RH) model*, $t_{j+\hat{x},j} = (-)^{j_y} + \delta t_x$, $t_{j+\hat{y},j} = 1 + \delta t_y$. Here $\delta\theta_{x,y}(j)$ and $\delta t_{x,y}$ are real random variables. Numerical results of the wave functions for these two models are qualitatively similar. The model (i) belongs to the unitary ensembles. The model (ii) preserves the time reversal symmetry and belongs to the orthogonal ensembles which is suitable for a comparison with the critical continuum theory.

In the absence of the randomness, the system is invariant under the translation and the Hamiltonian is written in the momentum space:

$$H_0 = \int_{-\pi}^{\pi} \frac{dk_x}{2\pi} \int_0^{2\pi} \frac{dk_y}{2\pi} \psi_k^\dagger \begin{pmatrix} \cos k_y & \cos k_x \\ \cos k_x & -\cos k_y \end{pmatrix} \psi_k \quad (2)$$

with $\psi_k^\dagger = (c_k^\dagger, c_{k+(0,\pi)}^\dagger)$. We see that the zero-energy state is formed by the fermions with momentum $\mathbf{k}_1 = (\pi/2, \pi/2)$ and $\mathbf{k}_2 = (-\pi/2, \pi/2)$. Expanding it near $\mathbf{k}_1$ and $\mathbf{k}_2$, we find in the continuum approximation, H_0 becomes

$$H_0 = 2i \int d^2x \Psi^\dagger \left[\begin{pmatrix} \sigma_1 & 0 \\ 0 & -\sigma_1 \end{pmatrix} \partial_x + \begin{pmatrix} \sigma_3 & 0 \\ 0 & \sigma_3 \end{pmatrix} \partial_y \right] \Psi \quad (3)$$

with $\Psi^\dagger = 1/(4\pi^2) \int d\delta k_x d\delta k_y e^{-i\delta k_x x - i\delta k_y y} (\psi_{k_1}^\dagger, \psi_{k_2}^\dagger)$ and $c_{j_x, j_y} = (i^{j_x+j_y}, i^{j_x-j_y}, i^{-j_x+j_y}, i^{-j_x-j_y}) \Psi$ where σ 's are the Pauli matrices. We assume that Ψ take the same value on four sites $(x, y) = (2m, 2n), (2m+1, 2n), (2m, 2n+1), (2m+1, 2n+1)$. Note that for weak impurities, the states near zero energy are described by the smooth function $\Psi(x)$. Therefore we can derive a continuum theory to describe those states near zero energy.

Before deriving the continuum theory with randomness, we first study the symmetries of the lattice Hamiltonian. Since the square lattice is bipartite, the transformation $c_{j_x, j_y} \rightarrow c'_{j_x, j_y} = (-1)^{j_x+j_y} c_{j_x, j_y}$ induces sign change of the Hamiltonian. It is expressed as $H(x) \rightarrow \gamma^\dagger H(x) \gamma = -H(x)$, $\gamma = \sigma_1 \otimes \sigma_1$ where $H = \int d^2x \Psi(x)^\dagger H(x) \Psi(x)$ is the continuum Hamiltonian. The contribution from the randomness $\delta H \equiv H - H_0$ has a form $\delta H = \int d^2x \Psi^\dagger g(x) \Psi$ if we ignore the

terms containing derivatives. The above condition implies that the most general form of the $g(x)$ is given by a linear combination of the 4 by 4 matrices: $\gamma_1 = \sigma_2 \otimes I$, $\gamma_2 = \sigma_1 \otimes \sigma_2$, $\gamma_3 = -\sigma_2 \otimes \sigma_1$, $\gamma_4 = I \otimes \sigma_2$ with coefficients $2a_i(x)$, $i = 1, \dots, 4$. The total Hamiltonian, after a change of basis to make $\gamma = -\sigma_3 \otimes I$, ($\Psi = T\Psi'$, $T = (I \otimes I + i\sigma_2 \otimes \sigma_1)/\sqrt{2}$), is given by

$$H = \int d^2x \Psi'^\dagger 2(I \otimes \sigma_1) \begin{pmatrix} 0 & D_+ \\ D_- & 0 \end{pmatrix} \Psi', \quad (4)$$

$$D_\pm = i[(\partial_x \mp a_1)\sigma_1 \mp (\partial_y \mp a_2)\sigma_3] \pm ia_3 \mp \sigma_2 a_4. \quad (5)$$

Note that the averaged Hamiltonian is invariant under translation $(x, y) \rightarrow (x+1, y)$. This requires that $\langle a_i \rangle = 0$ for $i = 1, 2, 3$. The averaged Hamiltonian is also invariant under reflection $(x, y) \rightarrow (x, -y)$. This implies that $\langle a_4 \rangle = 0$.

The zero-energy state of H satisfies

$$D_+ \psi_+ = 0, \quad D_- \psi_- = 0, \quad (6)$$

where $\begin{pmatrix} \psi_- \\ \psi_+ \end{pmatrix} = \Psi'$. Thus the random bond model contains two Dirac fermions in a continuum approximation, which is the famous fermion doubling of the lattice fermions. It causes several difficulties in numerical calculations. However, as we discuss below, *we can avoid the doubling as far as the $E = 0$ state is concerned*.

Now let us discuss about the doubling which arises from the lattice models. Since our model is a nearest neighbor hopping model on a bipartite lattice, we can write the Hamiltonian as

$$H = (\{c_+^\dagger\}, \{c_-^\dagger\}) \begin{pmatrix} O & \mathcal{D} \\ \mathcal{D}^\dagger & O \end{pmatrix} \begin{pmatrix} \{c_+\} \\ \{c_-\} \end{pmatrix} \quad (7)$$

where $\{c_+\} = \{c_{j_x, j_y} | j_x + j_y \text{ is even}\} = \{c_{++}, c_{--}\}$ is a set of fermion operators at one of the sublattices (+) and $\{c_-\} = \{c_{j_x, j_y} | j_x + j_y \text{ is odd}\} = \{c_{+-}, c_{-+}\}$ is the other (-). Here let us assume a system to be $L_x \times L_y$ with both odd L_x and L_y with a fixed boundary condition. (We label the sites as $(1, 1), \dots, (L_x, L_y)$.) Then a total number of sites $L_x L_y = N_+ + N_-$ is odd with $N_+ = N_- + 1$ where $N_+ = \#\{+\text{ sites}\}$, $N_- = \#\{-\text{ sites}\}$. Therefore $\mathcal{D}$ is a $N_+ \times N_- = N_+ \times (N_+ - 1)$ rectangular matrix. The Schrödinger equation is reduced to $\mathcal{D}\mathcal{D}^\dagger \phi_+ = E^2 \phi_+$ and $\mathcal{D}^\dagger \mathcal{D} \phi_- = E^2 \phi_-$. Any state with $E \neq 0$ has a pair state at $-E$ as ${}^t(\phi_+, \phi_-)_{-E} = {}^t(\phi_+, -\phi_-)_E$. Since $\det \begin{pmatrix} O & \mathcal{D} \\ \mathcal{D}^\dagger & O \end{pmatrix} = 0$ (note that $\mathcal{D}$ is a form of $N_+ \times N_+ - 1$), there is always zero-mode. The wave function at $E = 0$ satisfies

$$\mathcal{D}^\dagger \phi_+^{E=0} = 0, \quad \mathcal{D} \phi_-^{E=0} = 0. \quad (8)$$

Since $\dim \phi_+^{E=0} = N_+$, the first one gives $N_- = N_+ - 1$ equations for N_+ variables and there is a *one* dimensional non vanishing solution. (Therefore this zero-mode

is not degenerate in general.) On the other hand, since $\dim \phi_-^{E=0} = N_- = N_+ - 1$, the second one gives N_+ equations for $N_- = N_+ - 1$ variables, this is overdetermined. Therefore we have $\phi_-^{E=0} = 0$.

It gives an important restriction in the continuum model. It implies

$$\psi_- = 0 \quad (9)$$

in (6) since the transformation matrix between $|\Psi\rangle$'s and the site operators' $UT^{-1}U^{-1}$, only mixes operators within each sublattices. Therefore there is no degeneracy (doubling) for the $E = 0$ state. This $E = 0$ state is described by a usual two component spinor. Therefore the independent component is not four but two as far as the $E = 0$ state is concerned. We could avoid the doubling by using the above special conditions.

Random Gauge Model. — Now let us show numerical results for the random gauge model. In the previous numerical calculations for the π flux state with randomness [7], it was not possible to treat large two-dimensional systems. In the present work, however, we only need the eigenstate of a semi positive definite operator $\mathcal{D}\mathcal{D}^\dagger$ with zero eigen value. This and the fact that the $E = 0$ eigen states are generally not degenerate enable us to treat large size systems up to 801×801 . We take $\delta\theta_x(i)$ and $\delta\theta_y(i)$ to be uniform random numbers between $-V/2$ to $V/2$. In Fig. 1, the wave function is shown for $V = 0.2$ with 801×801 system where it is coarse grained over 4 sites on the plaquette. It seems to be neither localized nor extended in a usual manner. To understand the nature of the wave functions, the function $f(\alpha)$ obtained numerically are also shown. The box size dependence is carefully included in the calculations. [5] The maximum of $f(\alpha)$ gives a Hausdorff dimension which is always two here since the wave functions are on the square lattice. The value α_0 which gives the maximum of $f(\alpha)$ gives scaling of the dominant parts of the wave function. It is two for an extended states which gives uniform non-singular scaling. For a critical state, $\alpha_0 \neq 2$ and has a singular scaling. [8,9] In our case, it apparently deviates from 2 which means the zero-mode is critical.

Random Hopping Model. — The properties of random Dirac fermion models have been studied in detail lately. We study the random hopping model to make a comparison with the continuum theory. In the continuum approximation, the fact that $\delta t_{x,y}$'s are real further requires $\gamma g(x)\gamma = g(x)^*$. Thus we take with the coefficients $a_i(x)$, $i = 1, \dots, 4$ to be real. Note that $D_\pm$ happens to be the Hamiltonian of Dirac fermions with *imaginary* random gauge potentials (a_1, a_2) , an *imaginary* random chemical potential a_4 , and a real random mass a_3 . Let (a_1, a_2) , $2a_3$ and $2a_4$ to have a Gaussian distribution with width g_A , g_V and g_M respectively. Averaging over randomness generates four-fermion interactions with (g_A, g_V, g_M) as coupling constants. The one-loop beta functions for the above coupling constants is calculated in Ref. [3] as

$$\dot{g}_A = 32g_Mg_V \quad (10)$$

$$\dot{g}_V + \dot{g}_M = -8(g_V - g_M)^2 \quad (11)$$

$$\dot{g}_V - \dot{g}_M = -8(g_V + g_M)(g_V - g_M) - 8g_A(g_V - g_M) \quad (12)$$

(Note comparing to Ref. [3], here we have an *imaginary* random chemical potential, a real random mass and an *imaginary* random gauge potential. Thus the beta functions are modified accordingly with $(g_A, g_V, g_M) \rightarrow (-g_A, -g_V, g_M)$.) The renormalization-group (RG) equation has a fixed line $(g_A, g_V, g_M) = (g_A, 0, 0)$. Around the fixed line, $g_V - g_M$ is irrelevant (note $g_A > 0$) and thus we may set $g_V = g_M$. In this case $g_V + g_M$ does not flow, and non-zero $g_V + g_M$ will drive $g_A \rightarrow \infty$. Thus the fixed line has at least one (marginally) relevant direction. Two-fermion operators are also relevant. But those two-fermion operators do not appear since $\langle a_i \rangle = 0$, $i = 1, \dots, 4$ due to the discrete symmetries. Because of the *imaginary* random gauge potential, the results in Refs. [10], [3] imply that all charge neutral operators are irrelevant, except two-fermion operators and four-fermion operators discussed above. Thus the fixed line has only one marginally relevant direction. It suggests that the RH model have a line of critical points at zero energy which is described by the line of critical points in the random Dirac fermion model. Those results further suggest that, once we fix the energy to be zero, there is only one marginally relevant direction for the critical line. Thus the critical line in the RH model contains two relevant directions. Due to its relation to the random Dirac fermion model, the critical line in the RH model is exactly solvable, and critical exponents can be calculated exactly.

To describe the critical line, set $a_3 = a_4 = 0$. Let us assume the probability distribution of $(a_1(\mathbf{r}), a_2(\mathbf{r}))$ is given by $P \propto \exp(-\int d^2r \frac{1}{2g_A}[a_1^2(\mathbf{r}) + a_2^2(\mathbf{r})])$ where g_A , characterizing the strength of the gauge field fluctuations, is exactly marginal. Thus g_A can be used to parameterize different critical points on the critical line. In general critical exponents are functions of g_A .

Note that the energy E couples to an operator $\Psi_+^\dagger \Psi_+$. Since Ψ_+ and $\Psi_+^\dagger$ belong to two independent Dirac fermion models described by D_+ and D_- respectively, The scaling dimension of $\Psi_+^\dagger \Psi_+$ is simply the sum of the scaling dimensions of Ψ_+ and $\Psi_+^\dagger$. From Ref. [10], we see that the scaling dimensions of Ψ_+ and $\Psi_+^\dagger$ are equal to $1/2$, which implies that the scaling dimension of $\Psi_+^\dagger \Psi_+$ is 1. This means that the dynamics exponent $z = 1$ and the density of states scales as $N(E) \propto |E|$; both exponents are independent of g_A .

In the continuum theory (5), the zero-energy state can be solved exactly [1]. Introducing ϕ and χ through $a_\mu = \partial_\mu \phi + \epsilon_{\mu\nu} \partial_\nu \chi$. One can show that the zero-energy state (with, for example, $\gamma = 1$) is given by

$$\Psi_+ \propto e^{i\phi + \chi} \quad (13)$$

and the probability distribution of ϕ and χ is given by $P \propto \exp(-\int d^2r \frac{1}{2g_A}[(\partial\phi(\mathbf{r}))^2 + (\partial\chi(\mathbf{r}))^2])$. The wave function (13) with this distribution was studied in detail in Ref. [1] [2], and was shown to have a multifractal structure. The exact multifractal scaling function $f(\alpha)$ is defined on $d_- \leq \alpha \leq d_+$ and is given by [2]

$$f(\alpha) = 8 \frac{(d_+ - \alpha)(\alpha - d_-)}{(d_+ - d_-)^2} \quad (14)$$

where $d_{\pm} = 2(1 \pm \sqrt{\frac{g_A}{2\pi}})^2$ for $g_A < 2\pi$ and $d_+ = 8\sqrt{\frac{g_A}{2\pi}}$, $d_- = 0$ for $g_A > 2\pi$. We see that for weak randomness ($g_A < 2\pi$) $f(\alpha)$ is peaked at

$$\alpha_0 = 2 + \frac{g_A}{\pi} \quad (15)$$

The parameters $a_i(\mathbf{r})$ in the continuum model is directly related to the lattice randomness. We find that $a_1(j) = -\Lambda\delta t_{j,j+\hat{x}}(-)^{j_x}$, $a_2(j) = -\Lambda\delta t_{j,j+\hat{y}}(-)^{j_x+j_y}$, $a_3(j) = \Lambda\delta t_{j,j+\hat{x}}(-)^{j_x+j_y}$ and $a_4(j) = -\Lambda\delta t_{j,j+\hat{y}}(-)^{j_y}$ with a dimensional parameter Λ . Therefore if we chose δt_{ij} to satisfy $\delta t_{j,j+\hat{x}} = \delta t_{j+\hat{y},j+\hat{y}+\hat{x}}$ and $\delta t_{j,j+\hat{y}} = -\delta t_{j+\hat{x},j+\hat{x}+\hat{y}}$. Then $g_V = g_M = 0$ and the lattice model will be on the critical line. We used this parametrization and performed a numerical calculation on the above fine tuned RH model. We set $\delta t_{x,y}$ as Gaussian random numbers with a variance V^2 ($g_A = \Lambda^2 V^2$). In Fig. 2, we have plotted a singular scaling function $f(\alpha)$. In the calculation, g_A is obtained from a value of α_0 . The $f(\alpha)$ obtained from numerical calculation agrees well with the exact theoretical result Eq. (14).

In summary, we find that in the continuum approximation, the RH model at zero energy has a critical line with only one marginally relevant direction. This analytical result is supported by numerical calculations on the fine-tuned RH model. Therefore the RH model is described by the continuum random Dirac fermions as far as critical behavior is concerned. We also find that the zero-energy states of the general random bond models also appear to be critical according to our numerical results. However, the numerical calculations can not eliminate a possibility of localization with huge correlation length (more than 800 lattice spacings) which may appear in our problem due to the marginally relevant direction.

This work was supported in part by Grant-in-Aid from the Ministry of Education, Science and Culture of Japan (YH, XGW, MK) and NSF grants DMR-9411574 (XGW). XGW also acknowledges the support from A.P. Sloan Foundation.

-
- * electronic mail address: hatsugai@coral.t.u-tokyo.ac.jp
- [1] A. Ludwig, M. Fisher, R. Shankar and G. Grinstein, Phys. Rev. B **50**, 7526 (1994).
 - [2] C. C. Chamon, C. Mudry and X.G. Wen, preprint cond-mat/9510088.
 - [3] D. Bernard, preprint hep-th/9509137.
 - [4] T.C.Halsey, M. H. Jensen, L. Kadanoff, I. Procaccia, and B. Shraiman, Phys. Rev. A **33**, 1141 (1986).
 - [5] See H. Hiramoto and M. Kohmoto, Int. J. Mod. Phys. B **6**, 281 (1992) and references therein.
 - [6] Y. Hatsugai and M. Kohmoto, Phys. Rev. B **42**, 8282 (1990). J.H.Han, D.J.Thouless, H. Hiramoto, and M.Kohmoto, Phys. Rev. B **50**, 11365 (1994), Y. Hatsugai, M. Kohmoto and Y.-S. Wu, preprint cond-mat/9510171.
 - [7] Y. Hatsugai and P.A.Lee, Phys. Rev. B **48**, 4204 (1993).
 - [8] B. Huckenstein, Rev. Mod. Phys. **67**, 357 (1995).
 - [9] Y. Avishai, Y. Hatsugai and M. Kohmoto, Phys. Rev. B **51**, 13419 (1995).
 - [10] C.C. Chamon, C. Mudry and X.G. Wen, preprints cond-mat/9509054, cond-mat/9501066; A.A. Nersisyan, A. M. Tselik, and F. Wenger, Phys. Rev. Lett. **72**, 2628 (1994).

FIG. 1. The zero-mode wave function of the random gauge model and the corresponding $f(\alpha)$. The system size is 801×801 and $V = 0.2$. The errors by the finite size corrections in the box sizes is about the symbol sizes within the shown region but is larger in the other.

FIG. 2. Functions $f(\alpha)$'s for the zero-mode wave functions of the RH model: 401×401 and $V = 0.3$. The solid line is the analytical result. The different symbols are for the different randomness realizations.




Fig.1




Fig.2